

POST OPERATIVE INSTRUCTIONS

- LUBRICATING DROPS: Your eyes may burn and may feel scratchy for several days after surgery. Use lubricating eye drops as often as needed.
- PLASTIC SHIELD: You will receive a plastic shield on the day of surgery. Please wear the plastic shield while you are sleeping or lying down for <u>7 days/nights</u>.
- SUNGLASSES: Wear sunglasses (any pair with UV protection) whenever you are outdoors for <u>1 month.</u>
- RUBBING YOUR EYE: Do not put any pressure or rub your eye for <u>2 weeks</u>.
- MAKE-UP Avoid wearing make-up for <u>1 week</u>.
- CLEANING YOUR EYE: You should wipe off the drop residue gently with a clean tissue.
- EATING AFTER SURGERY: You may eat what you like after surgery but do not drink alcohol for <u>24 hours</u> as this may interact with the anesthesia medications you have had during surgery.
- ACTIVITES: You may resume light activities, but avoid heavy lifting, straining, or intense exercising for <u>2 weeks</u>. No swimming under water or diving for <u>1 month</u>. Golf or tennis is permissible after <u>1 week</u> with protective eye wear. Avoid dirty or dusty activities (e.g. gardening or home renovation) for <u>2 weeks</u>.
- SHOWERING/BATHING: You may take a bath anytime or shower from neck down but you must avoid getting soap or water in the operated eye <u>2 weeks</u>.
- DRIVING: You should avoid driving for the first <u>48-72 hours</u> following your surgery unless your surgeon tells you otherwise. Your vision may seem clearer but you still may not be legally allowed to drive.
- RESUMING WORK: For office or desk jobs resume work within <u>1-3 days</u>, for those who do more active work that requires lifting heavy objects and/or working in dusty environments we suggest <u>2 weeks</u> off work.
- EXTENDED TRAVELLING: Please refrain from booking a flight or travelling abroad for the first 14 days following your surgery.
- GLASSES: it is normal for the operated eye to feel blurry for several hours or even a few days. Often the vision clears up slowly but it can also be sharp within a day. Everyone is different and each eye will heal differently. You will see one of our doctors in <u>5 to 7 days</u>. You will see your optometrist <u>3-4 weeks</u> after having your surgery.



DROPS INSTRUCTIONS

	Vigamox (Moxifloxacin) ✔	Maxidex ↓	Prolensa or Bromfenac ✔
On the day of surgery Start when you go home	1 drop every 2 hours	1 drop every 2 hours	None
Starting the day after surgery Use drops until bottle is empty	Continue 1 drop/ 3 times a day until the drops run out	Continue 1 drop/ 3 times a day until the drops run out	Continue 1 drop a day until the drops run out

- ✓ Wait 1-2 minutes between all drops.
- ✓ You CAN use **LUBICATING** drops while using medication drops.
- ✓ KEEP using your **GLAUCOMA** drops if you are prescribed any.
- ✓ <u>There is a **REFILL** on the medication prescription you received for your first surgery</u>. This refill is for your second eye surgery, if applicable.

EMERGENCY CALLS:

PLEASE CALL US IF YOU EXPERIENCE ANY OF THE FOLLOWING IN THE FIRST WEEK

Decreased vision Increased redness Increased pain Increased floaters

FIRST 10 DAYS

- DURING OFFICE HOURS (8am-4:30pm): CALL 416-792-3043 or 416-748-2020
- AFTER HOURS Please go to your nearest Emergency Department

AFTER 10 DAYS

- DURING OFFICE HOURS (8am-4:30pm) CALL: 416-792-3043 or 416-748-2020
- **AFTER HOURS** AND IT IS AN EMERGENCY, GO TO YOUR NEAREST EMERGENCY DEPARTMENT.

If you are admitted to a hospital within <u>10 days</u> of the procedure for any medical reason, please notify our office.